

AD _____

GRANT NUMBER DAMD17-95-1-5038

TITLE: HIV Vaccines Based on Novel MULV-HIV Fusion Proteins

PRINCIPAL INVESTIGATOR: Abraham Pinter, Ph.D.

CONTRACTING ORGANIZATION: Public Health Research Institute
New York, New York 10016

REPORT DATE: July 1998

TYPE OF REPORT: Annual

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

1998 1016056

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 1998	3. REPORT TYPE AND DATES COVERED Annual (26 Jun 97 - 25 Jun 98)	
4. TITLE AND SUBTITLE HIV Vaccines Based on Novel MULV-HIV Fusion Proteins			5. FUNDING NUMBERS DAMD17-95-1-5038	
6. AUTHOR(S) Abraham Pinter, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Public Health Research Institute New York, New York 10016			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) We have continued our efforts to develop an effective HIV-1 vaccine based on the V1/V2 domain of gp120. Studies with human sera and with antibodies isolated from immunized animals have shown that the V1/V2 domain contains highly conserved epitopes that can act as potent neutralizing targets for a broad range of primary HIV-1 isolates. We have prepared a prototype immunogen, based on the Case-A2 isolate, that contains the clade B V2 consensus sequence, and have shown that this antigen induces highly crossreactive anti-V1/V2 antibodies in both rats and macaques, and that specific fractions of these antibodies possess potent cross-neutralizing activity. We are continuing to analyze this immunogen in both the rat and macaque models, and we expect to further analyze the humoral responses to immunization with this protein, including the isolation of monoclonal antibodies against various epitopes in the V1/V2 domain. Our goal during the coming year is to more fully characterize the epitopes mediating the potent neutralization, and to modify the immunogen so that those epitopes are presented more specifically and efficiently. The macaque immunizations will provide sufficient amounts of antibodies for full characterization, and may allow challenge studies to be performed with appropriate SHIV strains.				
14. SUBJECT TERMS HIV Vaccines, Novel MULV-HIV, Fusion Proteins			15. NUMBER OF PAGES 12	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

✓
Where copyrighted material is quoted, permission has been obtained to use such material.

✓
Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

✓
Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

✓
In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

✓
For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


 8/6/98
PI - Signature Date

Table of Contents

Front Cover.....	1
SF 298, Report Documentation Page.....	2
Foreword.....	3
Table of Contents.....	4
Introduction.....	5
Body- Progress Report.....	5
Conclusions and Future Directions.....	10
Bibliography of publications and meetings abstracts.....	12

Introduction

Whereas it is known that some human immune sera possess potent neutralizing activities for primary viruses, the identity of the target epitopes mediating this neutralization is unknown, and currently available immunogens have not been able to induce such activities. Recent evidence from our laboratory suggests that the V1/V2 domain of HIV-1 gp120 contains epitopes that are potent neutralization targets for macrophage-tropic HIV-1 isolates. The objectives of this study were to elicit V1/V2-specific antibodies by immunization with a recombinant protein that expressed the isolated V1/V2 domain and to determine the breadth and potency of HIV-1 neutralization by these antibodies.

Progress Report: Results of macaque immunization studies

Whereas it is known that some human immune sera possess potent neutralizing activities for primary viruses, the identity of the target epitopes mediating this neutralization is unknown, and currently available immunogens have not been able to induce such activities. Recent evidence from our laboratory suggests that the V1/V2 domain of HIV-1 gp120 contains epitopes that are potent neutralization targets for macrophage-tropic HIV-1 isolates (●). The objectives of this study were to elicit V1/V2-specific antibodies by immunization with a recombinant protein that expressed the isolated V1/V2 domain and to determine the breadth and potency of HIV-1 neutralization by these antibodies.

The V1/V2 domain of a clinical HIV-1 isolate (Case-A2) was expressed as a fusion glycoprotein in CHO cells. This sequence was selected since the central portion of its V2 domain was almost identical to the clade B consensus sequence of this region (Wang et al., JVI 69, 2708-15, 1995). The carrier sequence chosen was derived from the N-terminal domain of the murine leukemia virus SU protein, gp70 (Kayman et al., JVI 68, 400-410, 1994). The Case-A2 V1/V2 protein was purified by affinity chromatography on a Ni-NTA column, utilizing a His6 tag incorporated near the N-terminus of the carrier gp70 sequence. The protein was shown by SDS-PAGE to be >90% pure.

Despite the apparent purity of the V1/V2 fusion protein, a radioimmunoprecipitation analysis of the labeled protein demonstrated the presence of two conformational forms of the antigen, which differed both in their reactivity with different monoclonal antibodies and their mobility on SDS gels when analyzed under nonreducing conditions (Fig 1). Both bands were recognized by K10A11, an antibody directed against a site in the gp70-derived carrier domain, and by C9B6, a mab directed against a linear V2 epitope (not shown). The upper band was recognized by K19B3, a mab directed against a conserved V1/V2 epitope, while the lower band was recognized by SC258 and 697D, mouse and human mabs directed against conformation V2 epitopes, as well as by several other mouse and human mabs directed against native conformational epitopes and by a number of human sera. This preferential recognition of lower band by the human antibodies suggested that it represented the correctly folded form, while the upper band represented an alternative, presumably nonnative conformation. The two bands coalesced after reduction of disulfide bonds with DTT, confirming that they represented distinct disulfide-bonded conformers. We were able to fractionate these forms by affinity chromatography on a column to which

19981016 056

monoclonal antibody SC258 was immobilized. The K19-reactive form was present in the flow through, while the native form was eluted by low pH buffer.




Fig 1 Detection of two distinct conformational forms of the Case-A2 V1/V2 fusion protein. 35S-cysteine labeled Case-A2 V1/V2 fusion protein was immunoprecipitated with the indicated monoclonal antibodies, and then deglycosylated with PNGase F before SDS-PAGE analysis. The PNGase F deglycosylation was necessary to remove glycan-dependent heterogeneity which resulted in broadened bands. Two bands differing by ~2kD in apparent molecular weight were resolved upon analysis under nonreducing conditions, while only a sharp single band with lower mobility was resolved after reduction of disulfide bonds with DTT.

Three rhesus macaques (# 6876, 6876 and 7026) were immunized with purified Case-A2 V1/V2 protein in the presence of Ribi RAS triple adjuvant (Monophosphoryl Lipid A {MPL}, Trehalose Dicorynomycolate {TDM}, and Cell Wall Skeleton {CWS} at initial doses of 25 µg/kg. Animal 6876 was boosted with the unfractionated antigen at 5 µg/kg, while the other two V1/V2-immunized animals (#7014 and #7026) were boosted with the SC238 Mab affinity-purified fraction of the V1/V2 protein. A fourth animal (#6874) was immunized with an equivalent amount of the gp70-related carrier sequence. The animals were bled prior to each immunization and at weekly intervals following each immunization. All three animals generated significant antibody titers against the fusion protein immediately after the first boost. These titers decayed after several months, and a potent anamnestic response was observed in all animals following the second boost.

Both monkeys immunized with the purified Case-A2-V1/V2 fusion protein produced antibodies that reacted with heterologous gp120s as well as with V1/V2 domains derived from *env* sequences of a number of unrelated HIV-1 isolates, including one Thai clade E sequence (Fig 2). These antibodies appeared to be recognizing common conserved sequences, as evidenced by the fact that almost all of the reactivity of antibodies induced by the Case-A2 V1/V2 sequence for two unrelated recombinant Env proteins (derived from the Ba-L and 451 isolates) was absorbed by




Fig 2 Crossreactivity of V1/V2-specific macaque sera (after absorption of gp70-reactive antibodies on p621 column) of with various V1/V2 fusion proteins, demonstrating crossreactivity with two distant V1/V2 sequences, including one from a Thai clade E isolate.

the heterologous SF162 V1/V2 protein (Fig. 3). In initial assays, we found that sera of the three animals immunized with the V1/V2 protein, but not that of the control animal, were able to neutralize the macrophage-tropic NL-HX-ADA virus. These results indicate that the Case-A2 V1/V2 fusion protein was able to induce crossreactive antibodies against native gp120 epitopes.




Fig. 3. Crossreactivity of V1/V2-specific macaque sera of with recombinant gp120s

To further characterize these antibodies, the V1/V2-specific IgG fraction of the immune sera was isolated by sequential immunoaffinity chromatography on a column containing the immobilized gp70-related carrier protein, followed by passage over a column containing the complete V1/V2 fusion protein (Fig. 4). This resulted in removal of >95% of all V1/V2-reactive antibodies. A portion of the V1/V2-specific antibodies bound to the second column were recovered by sequential elution with low pH buffers followed by elution with buffer containing 8M guanidine hydrochloride. After extensive buffer exchange, the isolated antibodies were quantitated and tested for HIV-1 neutralizing activities.





Fig. 4 Fractionation of V1/V2-specific antibodies present in sera of macaques immunized with the Case-A2 V1/V2 fusion protein.

All of the eluted antibody fractions possessed neutralizing activities for a number of macrophage-tropic isolates, with the lower pH and GuHCl antibody fraction generally being more potent than that eluted at pH 2.4 (Fig. 5). In addition to NL-HX-ADA, viruses neutralized included Ba-L, a recombinant derived from NL-HX-ADA that contained the Case-A2 V1/V2 domain, and a clade C primary isolate from Malawi, 92MW965C.

Macaque #6876 serum



Macaque #7026 serum





Fig 5. Neutralization of macrophage-tropic isolates by V1/V2-specific macaque sera

A different result was obtained for one primary isolate, US716B (Fig. 6). Whereas the 6876 serum and V1/V2-specific antibodies isolated from this serum also neutralized this virus, the 7026 serum enhanced infectivity. This enhancing activity was removed upon absorption of the anti-V1/V2 antibodies, and was recovered in the low pH eluates of this column, demonstrating that it was antigen-specific. Interestingly, the antibodies eluted in the GuHCl fraction of this column possessed neutralizing activity for the US716 isolate. This indicated that this serum possessed both neutralizing and enhancing antibodies for the US716B virus, and that the enhancing activity dominated in the unfractionated serum.

Neutralization of 92US716B by fractionated 6876 serum



Neutralization/enhancement of 92US716B by fractionated macaque 7026 serum




Fig 6. Neutralization/enhancement of primary HIV-1 isolate US716B by V1/V2-specific macaque sera

An unexpected result was that in contrast to the efficient neutralization of NL-HX-ADA and NL-Case-A2-ADA, these antibodies did not neutralize two related molecular recombinants that contained the identical V1/V2, but with T cell-tropic V3-V5 regions (Fig. 7). This result suggested that the neutralizing activity of these antibodies was specific for M-tropic isolates. This supports a model in which the key V1/V2 epitopes that are targeted by these antibodies function specifically in CCR5-dependent infections, either by virtue of a direct interaction with the CCR5 receptor, or as a result of a conformational structure that is specific for macrophage-tropic envelope proteins.




Fig. 7. Comparative neutralizations of macrophage-tropic and T cell-tropic HIV-1 isolates by V1/V2-specific macaque sera

These studies have demonstrated that our Case-A2 V1/V2 fusion protein is capable of inducing potent neutralizing antibodies against native epitopes in the V1/V2 domain. However, the yield of this class of antibodies in the resulting immune sera has been low, and the majority of antibodies produced appeared to be directed against epitopes that do not mediate neutralization and that may be carried on non-native forms of the immunogen. Our challenge is to learn how to modify the immunogen and/or immunization protocols so that the focus of the immune response is directed against the neutralizing targets. We are currently performing mutagenesis studies of our V1/V2 fusion protein to define the role of individual residues in determining folding, immunoreactivity and immunogenicity of this protein. Ideally, we would like to map the different classes of epitopes. This would allow us to hopefully modify the immunogen so that the deleterious epitopes are eliminated, while retaining and perhaps enhance the immunogenicity of the protective epitopes. The broad crossreactivity and potent cross-neutralizing activities we have been able to induce in rodents and macaques with our current form of the V1/V2 immunogen is very encouraging, and suggests that this vaccine, and improved versions derived from the current immunogen, may induce protective responses in humans as well.

Conclusions and Future Directions

These studies have confirmed the potency of antibodies against native epitopes in the V1/V2 domain, and have demonstrated that our Case-A2 V1/V2 fusion protein is capable of inducing such antibodies. However, the yield of this class of antibodies has been low, while the majority of the antibodies produced were directed against epitopes that do not mediate neutralization and that may be carried on non-native forms of the immunogen. In addition, we have found that a subfraction of the antibodies induced possesses enhancing activity against at least one primary isolate. Our challenge during the coming year is to define the relevant neutralizing and enhancing epitopes, and to learn how to modify the immunogen and/or immunization protocols so that the focus of the immune response is directed specifically towards the neutralizing targets.

During the coming year we shall extend these studies in the following directions.

- 1- We shall continue immunogenicity studies with our current V1/V2 prototype vaccine in rodents and macaque models. The rat studies will allow us to quickly evaluate different immunogen and adjuvant combinations, and to isolate additional monoclonal antibodies against specific V1/V2 epitopes. The macaque model has several advantages. First, it is important to determine whether primates exhibit a similar humoral response to this immunogen, before trials in humans can be considered. The macaque model will allow us to test this, and also allow the evaluation of the protective effects of these immune, using appropriate SHIV challenges. This animal model will allow the determination of the most effective adjuvants for inducing these antibody responses, and may also allow the determination of whether these vaccines induce CTL responses. Secondly, a difficulty in fully evaluating the activities of the immune rat sera is the limited amounts of sera that can be obtained from these animals. The macaques are much larger animals, and can therefore provide sufficient antisera for multiple analyses and fractionations, by ourselves as well as by independent laboratories.

We will repeat the studies described above in larger scale and in additional animals, to fully characterize the potency and breadth of neutralizing activity of the anti-V1/V2 antibodies induced by the Case-A2 V1/V2 immunogen, and to see how common such responses are. We will also evaluate the potency of additional V1/V2 immunogens, including other sequences isolated from macrophage-tropic clade B isolates, as well as newer constructs derived from non-clade B sequences and modified immunogens in which the gp70 sequences have been removed and replaced with appropriate T helper epitopes. Depending on the results of the epitope mapping studies designed to distinguish neutralizing from non-neutralizing epitopes, we may also be able to design and test second generation immunogens which have been optimized for the retention of neutralization epitopes

2- Additional future directions we intend to take this study include the improvement of these V1/V2 immunogens to focus the humoral responses towards the relevant epitopes that induce the most potent neutralizing responses. This will involve evaluation of alternate strategies to either remove or modify the gp70-derived fusion sequences, to avoid the generation of antibodies against the carrier sequences. More importantly, our studies have shown that the V1/V2 domain is an immunologically complex structure. The most potently neutralizing antibodies are directed against the conserved conformational epitopes, and these are the most relevant ones for protection. A second class of antibodies are directed against the T15K V2 linear epitope. Whereas these also neutralize many viruses, they are less potent. One possibility is that mutating this sequence may increase the immunogenicity of the conformational epitopes. Previous studies with monoclonal antibodies to V1/V2 have shown that most of these have little, if any neutralizing activity. Such antibodies, if present in excess may be deleterious, since they may block the interaction of the neutralizing antibodies, thereby inhibiting their activities. Finally, we have found in some human sera the presence of anti-V1/V2 antibodies that actually enhance the infection by some strains of HIV-1. Such antibodies could be harmful if induced in response to immunization.

We have initiated mutagenesis studies of our V1/V2 fusion protein to define the role of individual residues in determining folding, immunoreactivity and immunogenicity of this protein. Initial targets being addressed are individual N-linked glycosylation sites. We are also expressing smaller subdomains of the V1/V2 region, including proteins containing only the conserved stem of the V1/V2 domain and proteins in which the conserved stem sequences have been deleted. These studies should help us define the relevant neutralizing epitopes, and hopefully would allow us to eliminate deleterious epitopes, while retaining and perhaps enhance the immunogenicity of the protective epitopes. The broad crossreactivity and potent cross-neutralizing activities we have been able to induce in rodents and macaques with our current vaccine is very encouraging, and suggests that this vaccine, and improved versions derived from the current immunogen, may induce protective responses in humans as well.

e. Publications and Abstracts, 1997-8

S.H. Pincus, K.G. Messer, R. Cole, R. Ireland, T.C. Vancott, A. Pinter, D.H. Schwartz, B.S. Graham, G.J. Gorse and the AIDS Vaccine Clinical Trials Network. 1997. Vaccine-specific antibody responses induced by HIV-1 envelope subunit vaccines. *J. Immunol.*, 158, 3511-3520.

S.H. Pincus, K.S. Wise, W.J. Honnen, A. Pinter, B. Cole and R.L. Cole. 1998. Immunologic crossreaction between HIV p17 and *Mycoplasma Hyorhinis* variable lipoprotein. *AIDS Res. Hum. Retroviruses*, 14, 419-425.

A. Pinter, W.J. Honnen, S.C. Kayman, O. Troshev and Z. Wu. 1998. Potent neutralization of primary HIV-1 isolates by antibodies directed against epitopes present in the V1/V2 domain of HIV-1 gp120. Vaccine, In press.

Meeting Abstracts

G. Tan, W.J. Honnen, S.C. Kayman and A. Pinter. Development of sensitive GFP-based infectivity and neutralization assay for macrophage-tropic HIV-1 isolates that utilizes a continuous cell line expressing the co-receptor CCR-5. Submitted for the 9th Annual Meeting of the National Cooperative Vaccine Development Groups for AIDS, May 4-7, 1997, Bethesda, MD.

A. Pinter, W.J. Honnen, S.C. Kayman, O. Troshev and Z. Wu. Production of highly crossreactive antibodies with potent neutralizing activities for a broad range of primary HIV-1 isolates upon immunization of rodents with a fusion glycoprotein containing the native V1/V2 domain of gp120. Submitted for the 9th Annual Meeting of the National Cooperative Vaccine Development Groups for AIDS, May 4-7, 1997, Bethesda, MD.

A. Pinter, W.J. Honnen, S.C. Kayman, S. Reiken, G. Tan, O. Troshev and Z. Wu. The V1/V2 domain of gp120 contains broadly conserved epitopes that mediate potent cross-clade neutralization of primary HIV-1 isolates. 1997 ISV Symposium on Vaccinology, Leesburg VA, Sept. 8, 1997.

A. Pinter, M. Burkhart, W. Honnen, S. Kayman, S. Reiken, G. Tan, O. Trochev and Z. Wu. Characterization of Highly Conserved Epitopes in the V1/V2 Domain of HIV-1 gp120 that Mediate Potent Neutralization of Primary HIV-1 Isolates. Fifth Conference on Retroviruses and Opportunistic Infections, Chicago, IL., Feb. 1998.

A. Pinter, O. Trochev, W.J. Honnen, Z. Wu, S. Reiken, M. Lewis and S.C. Kayman. 1998. Potent neutralization of primary HIV-1 isolates by antisera induced upon immunization with a recombinant protein expressing the native V1/V2 domain of gp120. To be presented at the 12th World AIDS Conference, Geneva. July 1998.

O. Trochev, S. Kayman, W. Honnen, S. Reiken, M. Lewis and A. Pinter. 1998. Induction of potent humoral immune responses for primary macrophage-tropic isolates of HIV-1 in rhesus macaques immunized with a recombinant protein expressing the native gp120 V1/V2 domain. To be presented at the Institute of Human Virology Annual Meeting, August 1998.